

Ameliorative Potential of Chlorogenic Acid on Rotenone-Induced Neurotoxicity in *Drosophila Melanogaster* Model

Oluwatoyin Adenike Adeyemo-Salami^{1,*}, Opeyemi Jamiu Afonja¹, Olamuyiwa Faosiyat Adeleke¹, Adeola Oluwatosin Adedara², Amos Olalekan Abolaji²

¹Nutritional and Industrial Biochemistry Unit, Department of Biochemistry, College of Medicine, University of Ibadan, Ibadan, Oyo State, Nigeria.

²Drosophila Laboratory, Department of Biochemistry, College of Medicine, University of Ibadan, Ibadan, Oyo State, Nigeria.

Corresponding author:

Oluwatoyin Adenike Adeyemo-Salami, Nutritional and Industrial Biochemistry Unit, Department of Biochemistry, College of Medicine, University of Ibadan, Ibadan, Oyo State, Nigeria.

keywords:

Drosophila melanogaster, chlorogenic acid, rotenone, Parkinson's disease, oxidative stress, acetyl cholinesterase

Running Title:

Ameliorative potential of chlorogenic acid

Received: Oct 11, 2021

Accepted: Nov 14, 2021

Published: Nov 18, 2021

Editor: Wang Wang, Department of Pediatrics The University of North Carolina at Chapel Hill

DOI: 10.14302/issn.2641-4538.jphi-21-3993

Abstract

Chlorogenic acid (CA), abundantly found in green coffee beans, is a phenolic compound with antioxidant and anti-inflammatory properties

amongst others. Exposure to rotenone, a natural pesticide, induces Parkinsonism (a type of neurodegeneration) through the induction of mitochondria dysfunction and oxidative stress. Phytochemicals with antioxidant properties may be promising in attenuating this condition. In this research, the ameliorative role of CA on rotenone-induced toxicity in *Drosophila melanogaster* was evaluated.

Drosophila melanogaster (Harwich strain, 1 - 3 days old) was used. 6 groups of five vials each with 50 flies/vial were exposed to CA (0; control (2% ethanol), 7.5, 15, 30, 45 and 60 mg/kg diet) for 28 days in the longevity analysis. A 28-day survival assay was carried out with rotenone (0, 250 and 500 μ M). CA (30 mg/kg diet) was selected to evaluate its ameliorative potential on rotenone. For the study, the flies were divided into four groups of five vials each and exposed to CA and rotenone; Group A- control (2% ethanol), Group B- CA only, Group C- rotenone only and Group D- CA (30 mg/kg diet)+ rotenone (500 μ M) for 7 days. Thereafter, the homogenate was evaluated for oxidative stress status, rate of emergence, negative geotaxis and acetyl cholinesterase activity.

CA (30 mg/kg diet) extended the lifespan of flies by 21.4%. Also, CA ameliorated rotenone-induced perturbation in catalase, glutathione-S-transferase and acetylcholinesterase activities, total thiol and glutathione levels, and behavioral deficit ($p < 0.05$).

CA may have ameliorative effect against rotenone-induced toxicity and Parkinsonism.

Introduction

Chlorogenic acid (CA) is a polyphenol from the family of hydroxycinnamic acid. It is one of the main polyphenols in the human diet with many health-promoting properties. CA can be found in foods and herbs such as coffee beans, apples, tea, grapes, tomato etc. [1, 2]. Furthermore, it has been found that chlorogenic acid has antioxidant, anti-inflammatory, anticancer, antilipidemic, antidiabetic, antihypertensive and anti-neurodegenerative activities [3- 8].

Rotenone (ROT) is a commonly used natural pesticide from the roots of *Derris elliptica* and *Lonchocarpus utilis* [9]. It can cross the blood brain barrier, due to its lipophilic property and causes neurotoxicity [10, 11]. ROT blocks the mitochondrial electron transport chain through the inhibition of complex I which causes cytotoxicity due to altered redox cycling and oxidative stress [12]. It has been observed that high doses of ROT can induce generalized neurodegeneration, therefore, investigations have been carried out on low-dose systemic administration of this compound in rodents. In this condition, there is slow and specific loss of dopaminergic neurons and impaired mobility in animals treated with ROT, which are attributes of Parkinson's disease [13].

Physical aberrations characterized by Parkinson's disease (PD) (a type of neurodegenerative disease) include progressive impairment in mobility ability such as bradykinesia, tremor and rigidity. Studies have shown that PD resulted from the loss of dopaminergic neurons in substantia nigra pars compacta (SNpc) and the formation of intra neuronal proteinaceous inclusions called Lewy

bodies (LBs) in affected brain areas [14- 17]. Moreover, the pathology involves protein aggregates apart from LBs, various neurotransmitters and extensive regions of the nervous system [18].

In this study, using ROT-induced toxicity in *Drosophila melanogaster* as a model for neurodegeneration, the ameliorative effect of CA was evaluated.

Materials and Methods

Chemicals

All chemicals were of analytical grade. ROT and CA were procured from AK Scientific, 30023 Ahern Ave, Union City, CA 94587, United State of America.

Drosophila Melanogaster Culture

Drosophila melanogaster (Harwich strain) obtained from the National Species Stock Centre, Bowling Green, Oklahoma, U.S.A., were maintained and reared in the *Drosophila* Laboratory, Department of Biochemistry, University of Ibadan, Oyo State, Nigeria on cornmeal medium containing brewer's yeast (1% w/v), agar-agar (1% w/v), and nipagin (preservative, 0.08% v/w) at constant temperature (23 ± 2 °C) under 12 h dark/light cycle.

Rotenone and Chlorogenic Acid Exposure and Survival Rate Analyses

In order to determine the appropriate concentrations and duration of exposure to ROT and CA to be used for the main experiment, *D. melanogaster* (both genders, 1 to 3 days old) were allocated into three (3) groups of five vials with 50 flies each and administered ROT (0- control (2% ethanol), 250, and 500 μ M doses respectively) for 28 days for survival assay while CA was exposed to another six (6) groups of five vials with 50 flies each (0- control (2% ethanol), 7.5, 15, 30, 45, and 60 mg/kg diet respectively) for 28-day longevity assay. For both assays, daily mortality was recorded, and data were analyzed and plotted as percentage of live flies. The effect on antioxidant status of exposure of six (6) groups of five vials with 50 flies each to CA (0- control(2% ethanol), 7.5, 15, 30, 45,

and 60 mg/kg diet respectively) was determined by assaying in the homogenate for catalase (CAT), glutathione-S-transferase (GST), hydrogen peroxide (H₂O₂), total thiol and reduced glutathione (GSH) after 7 days. Based on these data, 30 mg/kg diet of CA as well as 500 µM dose of ROT were selected for the study and used to determine the effect on biochemical parameters; acetylcholinesterase (AChE) activity (a marker for neurotoxicity) and oxidative stress-antioxidant status ((H₂O₂, total thiol, GSH levels, and CAT and GST activities), as well as the rate of emergence of offspring and negative geotaxis.

Determination of Negative Geotaxis and Rate of Emergence of Offspring

Locomotor performance of CA and ROT-treated flies were investigated using the negative geotaxis assay [19, 20]. Emergence rate of *D. melanogaster* offspring after exposure and co-exposure to CA and ROT were evaluated as previously described by Farombi *et al.* [21].

Preparation of Sample for Biochemical Assays

For the determination of biochemical assays, 50 flies (of both gender)/ vial for 5 vials each per group, were exposed as described in each of the following groups: Group A (control)-ethanol (2%), Group B- CA only (30 mg/kg diet), Group C- ROT only (500 µM) and Group D- 500 µM ROT + 30 mg/kg diet CA for 7 days. At the end of the treatment period, flies were anaesthetized using CO₂, weighed, and homogenized in 0.1 M phosphate buffer, pH 7.0 (ratio of 1 mg:10 ml), and centrifuged at 4000 g for 10 min at 4°C in a Thermo Scientific Sorval Micro 17R centrifuge. Subsequently, the supernatants were separated from the pellets into labeled Eppendorf tubes and used for the determination of the activities of AChE, GST, CAT, and total thiol, H₂O₂ and GSH content. All the assays were carried out in duplicates for each of the five replicates.

Determination of Biochemical Parameters

Estimation of Protein Concentration

The concentration of protein was evaluated by the method of Lowry *et al.*[22].The sample (25 µl, 1:10

dilution) was mixed with 135 µl of distilled water, and added to 400 µl of alkaline copper sulphate reagent (Lowry reagent) followed by 15 minutes incubation time. Folin ciocalteau solution (1:5 dilution) was added and they were then incubated at room temperature for 20 mins. The absorbance was measured at a wavelength of 660 nm against blank and protein values were estimated from the standard curve of bovine serum albumin (BSA).

Estimation of Total Thiol Level

The method of Ellman [23] was used to determine total thiol content. The reaction mixture consists of 510 µl of 0.1 M phosphate buffer (pH 7.4), 20 µl of sample, 35 µl of 1 mM 5',5' – dithiobis- 2- nitrobenzoate (DTNB) and 35 µl of distilled water. The mixture was incubated for 30 mins at room temperature and the absorbance was measured at 412 nm.

Estimation of Hydrogen Peroxide Level

The level of H₂O₂ in the treatment was determined using the method of Wolff [24]. The reaction mixture consist of FOX 1 (10 ml of 100 mM xylenol orange, 50 ml of 250 mM ammonium ferrous sulfate, 10 ml of 100 mM sorbitol, 5 ml of 25 mM H₂SO₄ and 30 ml of distilled water) mixed with the sample. Incubation was carried out for 30 mins at room temperature and the absorbance was measured at 560 nm. The values, which were deduced from the standard curve, were expressed in micromole per milligram protein.

Estimation of Reduced Glutathione Level

The estimation of the level of GSH was carried out with the method of Jollow *et al.* [25] with little modifications. Sulphosalicylic acid (4%) in the ratio of 1:1 was used to precipitate the supernatant. The samples were incubated for 1 h at 4°C and then centrifuged at 5000 rpm for 10 mins at 4°C. The reaction mixture consists of 550 µl of 0.1 M phosphate buffer, 100 µl of supernatant and 100 µl of DTNB. The absorbance was read at 412 nm and the results were expressed as moles of GSH/gram tissue.

Estimation of Glutathione-S-Transferase Activity

The method of Habig and Jakoby [26] was used to assess the activity of GST. The reaction mixture comprised of 270 µl of a solution consisting of 20 µl of 0.25 M potassium phosphate buffer, pH 7.0, with 2.5 mM ethylene diamine tetra acetic acid (EDTA), 10.5 ml of distilled water and 500 µl of 0.1 M GSH at 25 °C, 10 µl of 25 mM 1-chloro-2,4-dinitrobenzene (CDNB; as substrate) and 20 µl of sample (1:5 dilution). The mixture was monitored for 5 min with 10 s intervals at 340 nm using a spectrophotometer.

Determination of Catalase Activity

CAT activity was assessed with the method of Aebi [27]. The reaction mixture consists of 50 mM of phosphate buffer (pH 7.0), 300 mM H₂O₂ and sample (1:50 dilution). The loss in absorbance of H₂O₂ was monitored for 2 min at 240 nm. The result was used to calculate CAT activity and expressed as µmol of H₂O₂ consumed per minutes per milligram of protein.

Determination of Acetyl Cholinesterase Activity

AChE activity was evaluated using the method of Ellman *et al.* [28]. The reaction mixture contained 135 µl of distilled water, 20 µl of 100 mM potassium phosphate buffer (pH 7.4), 20 µl of 10 mM DTNB, 5 µl of sample, and 20 µl of acetylthiocholine (8 mM). The reaction was monitored at a wavelength of 412 nm (for 5 min, 15 s intervals) in a SpectraMax microplate reader (Molecular Devices, USA). The data were expressed thereafter in µmol/min/mg protein.

Statistical Analysis

All of the experiments were replicated at least two times. The data are presented as the mean ± standard error of mean (SEM). To assess the significant differences among multiple groups under various treatments, One-way analysis of variance (ANOVA) was used. For all of the assays, differences with *p*-value < 0.05 were considered statistically significant.

Results

Longevity and Selected Biochemical Indices in Flies Exposed to Chlorogenic Acid

Life time exposure to CA (7.5, 30, 45 and 60 mg/kg diet) increased longevity of flies by 3.57%, 21.4%, 14.3% and 7.1% respectively while there was a reduction (-3.57%) at the 15 mg/kg diet (Fig. 1). In addition, CA maintained CAT (Figure 2A) and GST (Figure 2B) activities in most of the groups, and reduced H₂O₂ levels in the 15, 30 and 60 mg/kg diet (Figure 2C) after exposure to flies for 7 days when compared with the control flies (*p* < 0.05). CA (30, 45, and 60 mg/kg diet) concentrations maintained total thiol level in flies when compared with the control (Figure 2D). Moreover, 45 mg/kg diet of CA significantly increased GSH (Figure 2E) levels in flies when compared with the control group (*p* < 0.05). Generally, 30 mg/kg diet of CA prolonged longevity and maintained redox balance of the flies significantly (*p* < 0.05) when compared with the other groups and the control flies. Therefore, 30 mg/kg concentrations was selected to investigate the ameliorative role on ROT-induced toxicity in the flies.

Effects of Rotenone on Survival of *Drosophila Melanogaster*

Figure 3 shows the effects of ROT on survival of *D. melanogaster* after 28 days of treatment with 250 µM and 500 µM of rotenone concentrations. The group treated with 500 µM of ROT concentration shows an increase mortality rate when compared with the control (*p* < 0.05). Hence 500 µM concentration was chosen for the study.

Effects of Chlorogenic Acid on the Redox State of Rotenone Exposed Flies

Figure 4A shows the effects of CA and ROT on the H₂O₂ level in *Drosophila melanogaster* after 7 days of treatment. There was no significant effect upon administration of CA on the ROT-treated group when compared with control (*p* < 0.05). There was a significant elevation in CAT activity in the ROT-treated group which was ameliorated by CA (Figure 4B; *p* < 0.05). In addition, Figure 4C shows the levels of total thiol and GSH (Figure 4D), and GST activity (Figure 4E) in *D. melanogaster*




Figure 1. Influence of chlorogenic acid on longevity




Figure 2. Effect of chlorogenic acid on antioxidant status.

Values are expressed as mean ± standard error of mean of 50 flies/vial with 5 replicates per treatment group. CA- chlorogenic acid, *-significantly different from control at $p < 0.05$




Figure 3. Influence of rotenone on survival of *D. melanogaster* after 28 days of treatment

exposed to ROT and CA. CA significantly mitigated ROT-induced decrease ($p < 0.05$) in the levels of total thiol and GSH when compared with the control ($p < 0.05$). Also, CA significantly increased ROT-induced inhibition of GST activity to levels comparable with the control ($p < 0.05$). Comparing the CA treated ROT group with the ROT only group, administration of CA significantly ($p < 0.05$) reduced CAT activity (Figure 4B) and significantly increased total thiol level ($p < 0.05$) (Figure 4C).

Acetyl Cholinesterase and Behavioural Assays

A significant reduction in the climbing rate of flies was observed (Figure 5A, $p < 0.05$) in ROT-treated flies after 7 days when compared with the control. However, CA reversed ROT-induced behavioural deficit depicted by the climbing rate ($p < 0.05$). In addition, ROT had a significant adverse effect on the fertility of flies, however CA alleviated ROT-induced alteration in fertility depicted by eclosion rate after 7 days of treatment (Figure 5B, $p < 0.05$). CA also significantly ($p < 0.05$) alleviated ROT-induced inhibition of AChE activity to levels comparable to control (Figure 5C). Comparing the CA

treated ROT group with the ROT only group, administration of CA significantly ($p < 0.05$) increased the climbing and eclosion rates, and the AChE activity.

Discussion

In this study, it was demonstrated in *Drosophila melanogaster* that CA ameliorated the deleterious effects of ROT on AChE and GST activities, total thiol and GSH levels, and climbing and eclosion rates.

PD is an age-related neurodegenerative disease affecting more than 1% of the human population. It is the second most common neurodegenerative disorder with prevalence increasing with age, mostly among population above 60 years [29]. The symptoms of PD include progressive impairment in locomotive ability, tremor, rigidity, and bradykinesia [17]. Currently, a combination of genetic changes and environmental factors has been observed to be responsible for this condition, though controversially [30, 31]. Some of the environmental factors is the exposure to chemicals in pesticides and herbicides such as ROT, permethrin, MPTP (1-methyl 4-phenyl 1, 2, 3, 6-tetrahydropyridine), organochlorines,




Figure 4. Effect of chlorogenic acid on rotenone induced alterations on oxidative stress markers. Values are expressed as mean ± standard error of mean of 50 flies/vial with 5 replicates per treatment group. CA- chlorogenic acid, ROT- rotenone, *-significantly different from control flies at p < 0.05, #-significantly different from rotenone-treated flies at p < 0.05.




Figure 5. Influence of chlorogenic acid on negative geo taxis, flies' emergence and acetyl cholinesterase activity in rotenone-treated flies. Values are expressed as mean ± standard error of mean of 50 flies/vial with 5 replicates per treatment group. CA- chlorogenic acid, ROT- rotenone, *- significantly different from control flies at p < 0.05, #- significantly different from rotenone-treated flies at p < 0.05.

paraquat and 2, 4-dichlorophenoxyacetic acid [32-34]. In this study, ROT was used to induce Parkinsonism in Harwich strain of *Drosophila melanogaster*.

Currently, the modulatory role of several nutraceuticals to attenuate endogenous oxidative stress status has been considered as an effective approach to achieve neuroprotection [35- 37]. As stated earlier, CA is a phenolic compound found in food of plant origin which has anti-diabetic, antioxidant, anti-inflammatory, anticancer, antilipidemic, and anti-neurodegenerative properties [3-8]. Moreover, studies have shown that consumption of diets rich in antioxidant property contribute to extension of lifespan in an organism [38]. The fly-survival and longevity experiment based on CA showed that among the concentration range tested (7.5- 60 mg/kg diet), 30 mg/kg of CA was the most effective (Figure 1). Therefore, in regard to the ingestion of CA, moderate levels would be more beneficial to increase longevity. Moreover, CA significantly reduced the level of H₂O₂ with the 30 mg/kg diet and did not show any adverse effect on the activities of CAT and GST as well as the levels of total thiol and GSH (Figure 2).

In this study, it was observed that 28-day exposure of *D. melanogaster* to ROT elicit alterations in the levels of antioxidant enzymes activities (such as CAT and GST), perturbations in reduced total thiol and GSH levels, increased H₂O₂ level, induced behavioural deficit as well as decreased AChE activity. All the aforementioned detrimental effects observed (with the exception of H₂O₂ level) were mitigated by CA (Figures 4 and 5).

CAT catalyzes the conversion of H₂O₂ to water (H₂O) and oxygen. The most common one is a monofunctional heme-containing enzyme [39]. It is an intracellular antioxidant enzyme that protects biological macromolecules from oxidation by H₂O₂ [39]. This study shows an elevated CAT activity in flies exposed to ROT which was mitigated to levels comparable to control by CA (Figure 4B). This reveals the modulatory role of CA in obtaining an optimum level of H₂O₂ molecule in the cell, which is also essential in cellular signaling and this is

supported by the observation with the CA diets (Figure 2C).

H₂O₂ is a reactive oxygen species (ROS). Elevated levels of H₂O₂ indicate that a system is plagued by oxidative stress [39]. In this study it was observed that there was reduction in the level of H₂O₂ in flies treated with CA alone. However, ROT caused a significant accumulation of H₂O₂ which was mitigated, but not significantly, by CA (Figure 4A). Antioxidants play protective roles by scavenging reactive oxygen species and reducing the insults of oxidative damage [40]. Although CA did not significantly reduce H₂O₂ level in the co-administered flies, the reduction in CAT activity upon co-administration shows that CA has an ameliorative effect on ROT-induced toxicity but was not distinct enough on H₂O₂ level upon conclusion of the experiment.

Total thiols (protein and non-protein thiols) defend the system against oxidative damage [41]. GSH functions as part of the first line of antioxidant defense system in living organisms. In conjunction with GSH-dependent enzymes, it helps to block the propagation of free radical chain reactions and detoxify deleterious ROS [42]. GSH is an important antioxidant in living organisms that prevents oxidative damage to cellular components [42- 44]. Depletion of cytosolic GSH in neurons is one of the hallmarks in PD [45]. The decrease in levels of total thiols and GSH in the ROT challenged group evidenced from our study suggests a compromised defense system in *D. melanogaster*. However, this study shows that CA modulates ROT-induced reduction of thiol containing compounds, thus confirming its anti-oxidative capacity. Hence, CA may be an ameliorative agent against diseases associated with altered redox state of a system.

GST is a major phase II detoxification enzyme, which comprises of a large and multi-functional enzyme family primarily, involved in the detoxification of endogenous substrate by catalyzing the conjugation of the nucleophilic sulphur atom of GSH with a variety of reactive electrophilic exogenous compounds [34, 46, 47]. The result from our study shows that CA protected against

ROT-induced inhibition of GST activity. Since GST catalyses the conjugation of GSH with electrophilic molecules, GSH was rapidly used up in order to protect the flies. Therefore, the ameliorative mechanism of CA in ROT-induced toxicity may be by further enhancing activity of thiol-containing proteins like GST.

AChE is a vital enzyme of the cholinergic system that modulates learning, sleep and wakefulness, memory, attention, stress response and locomotor activities. It hydrolyzes acetylcholine to choline and acetate thereby terminating cholinergic neurotransmission between synapses [48, 49]. Our study revealed that CA significantly alleviated ROT-induced inhibition of AChE activity when compared with control. This finding suggests that the modulatory action of CA on AChE activity may improve cognitive impairment evident in neuro-related diseases.

In the negative geotaxis assay, it was observed that ROT decreased the climbing rate of the flies but the effect was mitigated in groups co-treated with CA. This shows that CA has an ameliorative potency against bradykinesia, which is one of the effects of ROT toxicity and a hallmark in PD. This is similar to the study conducted by Farombi *et al.* [21], in which kolaviron was able to modulate the effect of ROT in flies.

Flies' emergence was highly increased in the group treated with CA only but a barren effect was recorded in the group treated with ROT only as compared with the control group. Also, substantial flies' emergence was observed in the co-administered group treated. This shows that CA may have a highly ameliorative effect against reproduction deficit common in PD patients. This is in line with the study conducted by Namula *et al.* [50] in which CA, in addition with caffeic acid, was able to improve the quality of frozen-thawed boar sperm [50].

Conclusion

The present study demonstrates that chlorogenic acid showed substantial ameliorative potential on rotenone-induced Parkinsonism in *Drosophila melanogaster* as shown by the reversal of the detrimental effects of rotenone on the levels and activities of total

thiol, reduced glutathione, glutathione-S-transferase and acetylcholinesterase. The negative geotaxis and developmental assays further buttressed the ameliorative effect of chlorogenic acid. These findings indicate that chlorogenic acid might have therapeutic effects for Parkinson's disease. Hence, majorly consumed home diet should be fortified with chlorogenic acid or there should be an adequate intake of foods rich in chlorogenic acid to prevent or modulate the effect of rotenone exposure in our environment as well as delay or possibly halt the onset of neurodegenerative diseases such as Parkinsonism.

Data Availability Statement (DAS)

The data is available upon request from the Corresponding author.

Disclosure of Interest

The authors declare that there is no conflict of interest.

Abbreviations

CA- chlorogenic acid, ROT- rotenone, PD- Parkinson's disease, SNpc- substantia nigra pars compacta, LBs- Lewy bodies, AChE- acetylcholinesterase, DTNB- 5',5' - dithiobis- 2- nitrobenzoate, GST- glutathione - S - transferase, CAT-catalase, CDNB-1- chloro-2,4-dinitrobenzene, EDTA- ethylene diamine tetra acetic acid, GSH- reduced glutathione, ROS- reactive oxygen species, MPTP- 1-methyl 4-phenyl 1, 2, 3, 6-tetrahydropyridine.

References

1. Niggeweg R, Michael AJ, Martin C. Engineering plants with increased levels of the antioxidant chlorogenic acid. *Nat. Biotechnol.* 2004;22: 746–54.
2. Bhattacharyya S, Majhi S, Saha BP, Mukherjee PK. Chlorogenic acid- phospholipid complex improve protection against UVA induced oxidative stress. *J Photochem Photobiol B.* 2014;130: 293–8.
3. Yagasaki K, Miura Y, Okauchi R, Furuse T. Inhibitory effects of chlorogenic acid and its related compounds on the invasion of hepatoma cells in culture. *Cytotechnology* 2000; 33:229-35. DOI:10.1023/

- A:1008141918852.
4. Noratto G, Porter W, Byrne D, Cisneros-Zevallos L. Identifying peach and plum polyphenols with chemo preventive potential against estrogen-independent breast cancer cells. *J Agric Food Chem.* 2009; 57:5219-26. DOI:10.1021/jf900259m.
 5. Lee B, Choi HY, Jeong CH, Bu Y. Chlorogenic acid ameliorates brain damage and edema by inhibiting matrix metalloproteinase-2 and 9 in a rat model of focal cerebral ischemia. *Eur J Pharmacol.* 2012;689:89– 95.
 6. Hwang SJ, Kim YW, Park Y, Lee HJ, Kim KW. Anti-inflammatory effects of chlorogenic acid in lipopolysaccharide-stimulated RAW264.7 cells. *Inflamm Res.* 2014;63: 81-90. DOI: 10.1007/s00011-013-0674-4.
 7. Nabavi SF, Tejada S, Setzar WN, Gortzi O, Sureda A, Braidy N, et al. Chlorogenic acid and mental diseases: From Chemistry to Medicine. *Curr Neuropharmacol.* 2017;15:471- 9. DOI: 10.2174/1570159X1466616032 5120625.
 8. Yan Y, Zhou X, Guo K, Zhou F, Yang H. Use of chlorogenic acid against Diabetes mellitus and its complications, *J Immunol Res.* 2020; 2020:9680508. <https://doi.org/10.1155/2020/96805>.
 9. Soloway SB. Naturally occurring insecticides. *Environ Health Perspect.* 1976;14:109–17. <https://doi.org/10.1289/ehp.7614109>.
 10. Santiago RM, Barbieiro J, Lima MM, Dombrowski PA, Andreatini R, Vital MA. Depressive-like behaviors alterations induced by intranigral MPTP, 6-OHDA, LPS and rotenone models of Parkinson's disease are predominantly associated with serotonin and dopamine. *Prog Neuropsychopharmacol Biol Psychiatry.* 2010; 34:1104–14. <https://doi.org/10.1016/j.pnpbp.2010.06.004>.
 11. Swarnkar S, Singh S, Mathur R, Patro I, Nath C. A study to correlate rotenone induced biochemical changes and cerebral damage in brain areas with neuromuscular coordination in rats. *Toxicology* 2010; 272: 17–22. DOI:10.1016/j.tox.2010.03.019.
 12. Sherer TB, Betarbet R, Testa CM, Seo BB, Richardson JR, Kim JH, et al. Mechanism of toxicity in rotenone models of Parkinson's disease *J Neurosci.* 2003;23:10756-64. DOI:10.1523/JNEUROSCI.23-34-10756.2003.
 13. Fleming SM, Zhu C, Fernagut P, Mehta A, DiCarlo CD, Seaman RL, et al. Behavioral and immunohistochemical effects of chronic intravenous and subcutaneous infusions of varying doses of rotenone. *Exp. Neurol.* 2004;187:418–29.
 14. Forno L S. Neuropathology of Parkinson's disease. *J Neuropathol Exp Neurol.* 1996;55:259–72.
 15. Nussbaum RL, Polymeropoulos MH. Genetics of Parkinson's disease. *Hum Mol Genet.* 1997;6:1687–91. <https://doi.org/10.1093/hmg/6.10.1687>.
 16. Thomas B, Beal MF. Parkinson's disease. *Hum Mol Genet.* 2007;16:R183-94. <https://doi.org/10.1093/hmg/ddm159>.
 17. Jankovic J. Parkinson's disease: clinical features and diagnosis. *J Neurol Neurosurg Psychiatry.* 2008;79: 368–76.
 18. Kalia LV, Lang AE. Parkinson's disease. *Lancet (London, England)* 2015;386: 896–912. [https://doi.org/10.1016/S0140-6736\(14\)61393-3](https://doi.org/10.1016/S0140-6736(14)61393-3).
 19. Feany MB, Bender WW. A *Drosophila* model of Parkinson's disease. *Nature* 2000;404:394–8.
 20. Abolaji AO, Kamdem JP, Lugokenski TH, Farombi EO, Souza DO, da Silva Loreto EL, et al. Ovotoxicants 4-vinylcyclohexene 1,2-monoepoxide and 4- vinylcyclohexene diepoxide disrupt redox status and modify different electrophile sensitive target enzymes and genes in *Drosophila melanogaster*. *Redox Biol.* 2015;5:328–39.
 21. Farombi EO, Abolaji AO, Farombi TH, Oropo AS, Owa OA, Awunah MT. *Garcinia kola* seed flavonoid fraction (Kolaviron), increases longevity and attenuates rotenone-induced toxicity in *Drosophila melanogaster*.

- Pestic Biochem Phys. 2018;145:39–45. <https://doi.org/10.1016/j.pestbp.2018.01.002>.
22. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 1951;193: 265–75.
 23. Ellman GL. Tissue sulfhydryl groups. *Arch. Biochem. Biophys.* 1959;82: 70–7.
 24. Wolff SP. Ferrous ion oxidation in presence of ferric ion indicator xylenol orange for measurement of hydroperoxides. *Methods Enzymol.* 1994;233: 182–9.
 25. Jollow DJ, Mitchell JR, Zampaglione N, Gillette JR. Bromobenzene-induced liver necrosis. Protective role of glutathione and evidence for 3,4-bromobenzene oxide as the hepatotoxic metabolite. *Pharmacology* 1974;11: 151-69.
 26. Habig WH, Jakoby WB. Assays for differentiation of glutathione-S-transferases. *Methods Enzymol.* 1981;77: 398–405.
 27. Aebi H. Catalase *in vitro*. *Methods Enzymol.* 1984;105:121–126.
 28. Ellman GL, Courtney KD, Andres V, Feathers-Stone RM. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem Pharmacol.* 1961;7: 88–95.
 29. Reeve A, Simcox E, Turnbull D. Ageing and Parkinson's disease: Why is advancing age the biggest risk factor. *Ageing Res Rev.* 2014;14:19-30. DOI: 10.1016/j.arr.2014.01.004.
 30. Inamdar NN, Arulmozhi DK, Tandon A, Bodhankar SL. Parkinson's disease: Genetics and beyond. *Curr Neuropharmacol.* 2007;5: 99-113.
 31. Ball N, Teo WP, Chandra S, Chapman J. Parkinson's disease and the environment. *Front Neurol.* 2019 [cited 2021 April 27]. Available from: <https://doi.org/10.3389/fneur.2019.00218>.
 32. Brown TP, Rumsby PC, Capleton AC, Rushton L, Levy LS. Pesticides and Parkinson's disease- Is there a link?. *Environ Health Perspect.* 2006;114:156-64.
 33. Tanner CM, Kamel F, Ross GW, Hoppin JA, Goldman SM, Korell M. et al. Rotenone, paraquat and Parkinson's disease. *Environ Health Perspect.* 2011;119:866-72. DOI: 10.1289/ehp.1002839.
 34. Abolaji AO, Adedara AO, Adie MA, Vicente-Crespo M, Farombi EO. Resveratrol prolongs lifespan and improves 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced oxidative damage and behavioural deficits in *Drosophila melanogaster*. *Biochem Biophys Res Commun.* 2018;503:1042–1048. <https://doi.org/10.1016/j.bbrc.2018.06.114>.
 35. Jimenez-Del-Rio M, Guzman-Martinez C, Velez-Pardo C. The effects of polyphenols on survival and locomotor activity in *Drosophila melanogaster* exposed to iron and paraquat. *Neurochem Res.* 2010;35: 227–38.
 36. Dumont M, Beal MF. Neuroprotective strategies involving ROS in Alzheimer Disease. *Free Radic Biol Med.* 2011;51:1014-26.
 37. Prasad SN, Muralidhara. Evidence of acrylamide induced oxidative stress and neurotoxicity in *Drosophila melanogaster* - its amelioration with spice active enrichment: relevance to neuropathy. *Neurotoxicology* 2012;33:1254–64.
 38. Kumar A, Christian PK, Panchal K, Guruprasad BR, Tiwari AK. Supplementation of Spirulina (*Arthrospira platensis*) improves lifespan and locomotor activity in Paraquat-Sensitive DJ-1 β^{A93} flies, a Parkinson's Disease model in *Drosophila melanogaster*. *J Diet Suppl.* 2017;14:573–88.
 39. Nandi A, Yan LJ, Jana CK, Das N. Role of catalase in oxidative stress- and Age-associated degenerative diseases *Oxid Med Cell Longev.* 2019; 2019: 9613090. <https://doi.org/10.1155/2019/9613090>.
 40. Kurutas EB. The importance of antioxidants which play the role in cellular response against oxidative/nitrosative stress: Current state. *Nutr J.* 2016;15:71. DOI: 10.1186/s12937-016-0186-5.
 41. Ulrich K, Jakob U. The role of thiols in antioxidant

- systems. *Free Radic Biol Med.* 2019;140: 14- 27. DOI:10.1016/j.freeradbiomed.2019.05.035.
42. Lushchak V.I. Glutathione homeostasis and functions: Potential targets for medical interventions. *J Amino acids.* 2012; 2012: 736837. DOI:10.1155/2012/736837.
43. Sharma P, Jha AB, Dubey RS, Pessarakli M. Reactive oxygen species, oxidative damage and antioxidative defense mechanism in plants under stressful conditions. *J Bot.* 2012;2012: 217037. <https://doi.org/10.1155/2012/217037>.
44. Biller JD, Takahashi LS. Oxidative stress and fish immune system: phagocytosis and leukocyte activity burst. *An. Acad. Bras. Ciênc.* 2018 [cited 2021 April 30]. Available from: <https://doi.org/10.1590/0001-3765201820170730>.
45. Garrido M, Tereshchenko Y, Zhevtsova Z, Taschenberger G, Bahr M, Kugler S. Glutathione depletion and overproduction both initiated degeneration of nigral dopaminergic neurons. *Acta Neuropathol.* 2011;121:475–85.
46. Mathew N, Kalyanasundaram M, Balaraman K. Glutathione-S- Transferase (GST) inhibitors. *Expert Opin Ther Pat* 2006;16: 431-44. DOI:10.1517/13543776.16.4.431.
47. Suthar PC. Glutathione-S-Transferases: A brief on classification and GSTM1-T1 activity. *IJPSR* 2017;8:1023-7.
48. Ferreira-Vieira TH, Guimaraes IM, Silva FR, Ribeiro FM. Alzheimer's disease: Targeting the cholinergic system. *Curr Neuropharmacol* 2016;14:101-15. DOI: 10.2174/1570159X13666150716165726.
49. Trang A, Khandhar PB. Physiology, Acetyl cholinesterase. StatPearls Publishing (Internet): U.S.A. 2021 [cited 2021 April 28] Available from: <https://www.ncbi.nlm.nih.gov/books/NBK539735/>.
50. Namula Z, Hirata M, Wittayarat M, Tanihara F, Nguyen NT, Hirano T, et al. Effects of chlorogenic acid and caffeic acid on the quality of frozen-thawed boar sperm. *Reprod Domest Anim.* 2018;53:1600-4. DOI: 10.1111/rda.13288.