

# The Balance $2 \cdot f(O) - f(H)$ as a Cornerstone in Formulation of Electrolytic Systems

Anna M. Michałowska-Kaczmarczyk<sup>1</sup>, Tadeusz Michałowski<sup>2,\*</sup>

<sup>1</sup>Department of Oncology, The University Hospital in Cracow, Cracow, Poland.

<sup>2</sup>Department of Analytical Chemistry, Technical University of Cracow, Cracow, Poland.

## Abstract

The Generalized Electron Balance (GEB) concept, related to electrolytic redox systems, is considered according to principles of Generalized Approach to Electrolytic Systems (GATES). Two equivalent Approaches (I and II) to GEB are presented. The Approach I, when perceived in convention of the card game, is based on the common pool of electrons as money, introduced by electron-active elements, named as players; electron-non-active elements are called there as fans. The GEB obtained according to Approach II results from the linear combination  $f_{12} = 2 \cdot f(O) - f(H)$  of elemental balances:  $f_1 = f(H)$  for H, and  $f_2 = f(O)$  for O. Very important properties of  $f_{12}$  are presented here in details, and illustrated by a redox system where comproportionation reactions occur.

**Corresponding Author:** Tadeusz Michałowski, Department of Analytical Chemistry, Technical University of Cracow, Cracow, Poland, Email: [michalot@o2.pl](mailto:michalot@o2.pl)

**Citation:** Anna M. Michałowska-Kaczmarczyk, Tadeusz Michałowski (2018) The Balance  $2 \cdot f(O) - f(H)$  as a Cornerstone in Formulation of Electrolytic Systems. Journal of new Developments in Chemistry - 2(1):1-13. <https://doi.org/10.14302/issn.2377-2549.jndc-18-2177>

**Keywords:** Thermodynamics; Electrolytic systems; Redox titrations; Oxidation numbers.

**Received:** June 26, 2018

**Accepted:** Aug 06, 2018

**Published:** Sep 11, 2018

**Editor:** Dr. Ashish Kumar, Associate Professor and HOD - Department Of Chemistry, Lovely Professional University Phagwara, India.

## Introduction

The linear combination  $f_{12} = 2 \cdot f(O) - f(H)$  of elemental balances:  $f_1 = f(H)$  for H, and  $f_2 = f(O)$  for O, can be formulated for electrolytic redox and non-redox systems, of any degree of complexity, where charge ( $f_0 = \text{ChB}$ ) and elemental/core balances  $f_k = f(Y_k)$  for  $Y_k \neq H, O$  ( $k \geq 3, \dots, K$ ) are also formulated, according to Generalized approach to redox systems (GATES) principles. The set of  $K$  independent balances  $f_0, f_{12}, f_3, \dots, f_k$  is necessary for thermodynamic resolution of a redox system; all the balances are presented here in the form  $f_k = 0$  ( $k = 0, \dots, K$ ). A non-redox system is resolvable with use of  $K-1$  independent balances  $f_0, f_3, \dots, f_k$ . In a non-redox system,  $f_{12}$  is the dependent balance, indicated by the linear combination  $LC = \sum_{k=1}^K d_k \cdot f_k - f_0$ , where  $d_k$  are the oxidation numbers (ONs) for  $Y_1 = H, Y_2 = O$ , and  $Y_k$  ( $k \geq 3, \dots, K$ ).  $f_{12} = \sum_{k=3}^K d_k \cdot f_k - f_0$ . For a non-redox system, the LC is expressed by identity,  $0 = 0$ , i.e.,  $f_{12}$  is linearly dependent on  $f_0, f_3, \dots, f_k$ , and then For a redox system, LC is expressed by equation different from the identity, i.e.,  $f_{12}$  is the new balance in the redox system. Therefore, the linear dependency/independency of  $f_{12}$  from  $f_0, f_3, \dots, f_k$  is the general criterion distinguishing between non-redox and redox systems, of any degree of complexity.

The Approach II to GEB (formulated 2005) was preceded by the Approach I to GEB (formulated 1992); both approaches are equivalent, i.e.,

Approach I to GEB  $\Leftrightarrow$  Approach II to GEB (1)

Formulation of GEB according to Approach I can be perceived in the convention of the card game, based on the common pool of electrons as money, introduced by electron-active elements, named as players; electron-non-active elements are called there as fans. In a redox system, the arranged set  $(f_1, f_2, \dots, f_{k^*}, f_{k^*+1}, \dots, f_k)$  of  $K$  balances  $f_1, \dots, f_k$  involves  $K^*$  balances  $f_1, f_2, \dots, f_{k^*}$  for fans, and  $K - K^*$  balances  $f_{k^*+1}, \dots, f_k$  for players. In a non-redox system we have  $K^* = K$  fans, i.e., the number of players is  $K - K^* = 0$  in there; H and O are considered here as fans.

The prior knowledge of ONs for all elements of the system in question is not needed in the Approach II to GEB, contrary to the formulation of GEB according to

Approach I, where the knowledge of ONs for players is required.

All the regularities indicated above will be confirmed in an example presented below, and illustrated graphically. Further generalizing comments will also be presented.

## Redox Systems in Chemistry

In chemical laboratories, redox reactions are usually referred to processes occurred during redox titrations. Since redox reactions are so important class of chemical reactions, we should know their physicochemical nature thoroughly, from the titrimetric viewpoint.

Titration is considered as a kind of dynamic process, where  $V$  mL of titrant T is added, from the start, up to a given point of the titration, into  $V_0$  mL of titrand D, and  $V_0 + V$  mL of D+T mixture is obtained, if the volumes additivity is valid/tolerable. The species formed in the D+T systems are involved in the related balances.

In aqueous media, the species  $X_i^{z_i}$  exist as hydrates  $X_i^{z_i} \cdot n_{iW}$ ;  $z_i = 0, \pm 1, \pm 2, \dots$  is a charge, expressed in terms of elementary charge units,  $e = F/N_A$  ( $F = 96485 \text{ C} \cdot \text{mol}^{-1}$  – Faraday's constant,  $N_A = 6.022 \cdot 10^{23} \text{ mol}^{-1}$  – Avogadro's number),  $n_i = n_{iW} \geq 0$  is a mean number of water ( $W = \text{H}_2\text{O}$ ) molecules attached to  $X_i^{z_i}$ ; the case  $n_{iW} = 0$  is then also admitted. For ordering purposes we assume:  $X_2^{z_2} \cdot n_{2W} = \text{H}^{+1} \cdot n_{2W}$ ,  $= X_3^{z_3} \cdot n_{3W} \text{OH}^{-1} \cdot n_{3W}$ , i.e.,  $z_2 = +1$ ,  $z_3 = -1, \dots$ . Molar concentration of the species  $X_i^{z_i} \cdot n_{iW}$  is denoted as  $[X_i^{z_i}]$ ; for a static system (volume  $V_0$  mL), we have

$$[X_i^{z_i}] = 10^3 \cdot (N_i/N_A)/V_0 \quad (2)$$

and for a dynamic D+T system

$$[X_i^{z_i}] = 10^3 \cdot (N_i/N_A)/(V_0+V). \quad (2a)$$

It is advisable to start the balancing from the interrelations between numbers of particular entities:  $N_{0j}$  for components represented by molecules composing D and T, and  $N_i$  – for species  $X_i^{z_i} \cdot n_{iW}$  (ions and molecules) of  $i$ -th kind ( $i = 1, \dots, I$ ), where  $I$  is the number of kinds of the species in the D+T. The mono- or two-phase electrolytic D+T system thus obtained involves  $N_1$  molecules of  $\text{H}_2\text{O}$  and  $N_i$  species of  $i$ -th kind,  $X_i^{z_i} \cdot n_{iW}$  ( $i = 2, 3, \dots, I$ ), specified

briefly as  $X_i^{z_i} (N_i, n_i)$ , where  $n_i \equiv n_{iW} \equiv n_i H_2O$  is the mean number of hydrating water molecules ( $W=H_2O$ ) attached to  $X_i^{z_i}$ . The net charge of  $X_i^{z_i} \cdot n_{iW}$  equals to the charge of  $X_i^{z_i}$ :  $z_i + n_{iW} \cdot 0 = z_i$ . For ordering purposes, we write the sequence:  $H^{+1} (N_2, n_2)$ ,  $OH^{-1} (N_3, n_3)$ , ... , i.e.,  $z_2 = 1, z_3 = -1, \dots$ .

In the notation applied here,  $N_{0j}$  ( $j = 1, 2, \dots, J$ ) is the number of molecules of the component of  $j$ -th kind, including water, forming  $D$  in static ( $V=0$ ), or  $D$  and  $T$  in a dynamic  $D+T$  system. The  $D+T$  system thus obtained involves  $N_1$  molecules of  $H_2O$  ( $X_1^{z_1} = H_2O, z_1 = 0$ ) and  $N_i$  species of  $i$ -th kind,  $X_i^{z_i} \cdot n_{iW}$  ( $i = 2, 3, \dots, I$ ), denoted briefly as  $X_i^{z_i} (N_i, n_i)$ , where  $n_i \equiv n_{iW} \equiv n_i H_2O$ . The  $n_i = n_{iW}$  values are virtually unknown – even for  $X_2^{z_2} = H^{+1} [1]$  in aqueous media, and depend on ionic strength ( $I$ ) of the solution.

Electrolytic (redox and non-redox) systems are formulated according to the GATES principles [2-40]. For this purpose, the set of  $K+1$  balances  $f_0, f_1, f_2, f_3, \dots, f_k$  is obtained. The charged/ionic species of the system are involved in the charge balance

$$f_0 = \sum_{i=2}^I z_i \cdot N_i = 0 \quad (3)$$

$$\sum_{i=2}^I z_i \cdot [X_i^{z_i}] = 0 \quad (3a)$$

applicable for static and dynamic systems. Note that  $0 \cdot [X_m] = 0$  for a species  $X_m \cdot n_{mW}$  with zero charge ( $z_m=0$ ), e.g.,  $0 \cdot [H_2O] = 0$  ( $z_1=0$ ).

Free water particles ( $N_1$ ), and water bound in the hydrates  $X_i^{z_i} \cdot n_{iW}$ , are included in the balances:  $f_1 = f(H)$  and  $f_2 = f(O)$ :

$$f_1 = f(H) = 2N_1 + \sum_{i=2}^I (a_{1i} + 2n_{iW}) \cdot N_i - \sum_{j=1}^J b_{1j} \cdot N_{0j} = 0 \quad (4)$$

$$f_2 = f(O) = N_1 + \sum_{i=2}^I (a_{2i} + n_{iW}) \cdot N_i - \sum_{j=1}^J b_{2j} \cdot N_{0j} = 0 \quad (5)$$

Next, the linear combination

$$f_{12} = 2 \cdot f_2 - f_1 = \sum_{i=2}^I (2 \cdot a_{2i} - a_{1i}) \cdot N_i - \sum_{j=1}^J (2 \cdot b_{2j} - b_{1j}) \cdot N_{0j} = 0 \quad (6)$$

is obtained, where  $a_{ki}$  and  $b_{kj}$  in equations 4-6 are the numbers of atoms/cores  $Y_k$  ( $k = 1, \dots, K$ ) in  $X_i^{z_i} \cdot n_{iW}$ , and in the  $j$ -th component of the system, resp. The core

is perceived as a non-transformable cluster of atoms, of the same elemental composition (expressed by chemical formula) and structure; e.g. the set of species:  $HSO_4^{-1}$ ,  $SO_4^{-2}$  and  $FeSO_4$  contains  $SO_4^{-2}$  as the core. One can notice that: (a)  $N_1$  involved with  $H_2O$  as the species, (b) all  $N_{0i}$  related to water as the component of the system, and (c) all  $n_i = n_{iW}$  specified in  $f(H)$  and  $f(O)$  are cancelled within  $2 \cdot f(O) - f(H)$  in aqueous media. The necessity of prior knowledge of  $n_{iW}$  values in the balancing procedure is thus avoided. All these regularities are valid for non-redox and redox systems.

The elemental/core balances:  $f_3, \dots, f_k$ , interrelating the numbers of atoms/cores  $Y_k \neq H, O$  in components and species, are as follows

$$f_k = f(Y_k) = \sum_{i=1}^I a_{ki} \cdot N_i - \sum_{j=1}^J b_{kj} \cdot N_{0j} = 0 \quad (7)$$

( $k = 3, \dots, K$ )

All the balances  $f_0, \dots, f_k$  are presented here areas equations  $f_k = 0$ , see equations 3-7.

In further parts of this paper, we concern on the procedure known in elementary algebra as linear combination [19,41], in accordance with Noether's conceptual approach to linear algebra [42,43]. For this purpose, we consider first an example of titration in a redox system, where comproportionation [44] reactions occur. The conclusions arising in the formulation of this system (denoted as the system  $S$ , for brevity) will then be generalized later in discussion.

### System S: $KBrO_3 (C) \Rightarrow NaBr (C_0) + H_2SO_4 (C_{01})$

Let us consider the system, where  $V_0$  mL of  $D$  is composed of  $NaBr$  ( $N_{01}$  molecules) +  $H_2SO_4$  ( $N_{02}$  molecules) +  $H_2O$  ( $N_{03}$  molecules), and  $V$  mL of  $T$  is composed of  $KBrO_3$  ( $N_{04}$  molecules) +  $H_2O$  ( $N_{05}$  molecules). The  $D+T$  mixture thus formed involves the following species:

$H_2O$  ( $N_1$ ),  $H^{+1}$  ( $N_2, n_2$ ),  $OH^{-1}$  ( $N_3, n_3$ ),  $HBrO_3$  ( $N_4, n_4$ ),  $BrO_3^{-1}$  ( $N_5, n_5$ ),  $HBrO$  ( $N_6, n_6$ ),  $BrO^{-1}$  ( $N_7, n_7$ ),

$Br_2$  ( $N_8, n_8$ ),  $Br_3^{-1}$  ( $N_9, n_9$ ),  $Br^{-1}$  ( $N_{10}, n_{10}$ ),  $Na^{+1}$  ( $N_{11}, n_{11}$ ),  $K^{+1}$  ( $N_{12}, n_{12}$ ),

$HSO_4^{-1}$  ( $N_{13}, n_{13}$ ),  $SO_4^{-2}$  ( $N_{14}, n_{14}$ ).

### The Approach II to the system S

Denoting  $C_0 V_0 = 10^3 \cdot N_{01} / N_A$ ,  $C_{01} V_0 = 10^3 \cdot N_{02} / N_A$

CV = 10<sup>3</sup>·N<sub>04</sub>/N<sub>A</sub>, we formulate the fraction titrated [16,25,45]

$$\Phi = \frac{C \cdot V}{C_0 \cdot V_0} \quad (8)$$

It provides a kind of normalization in the related graphs, i.e., independency on V<sub>0</sub> value. The balances are as follows:

$$f_0 = \text{ChB} :$$

$$N_2 - N_3 - N_5 - N_7 - N_9 - N_{10} + N_{11} + N_{12} -$$

$$N_{13} - 2N_{14} = 0 \quad \Rightarrow \quad (9)$$

$$[H^{+1}] - [OH^{-1}] - [BrO_3^{-1}] - [BrO^{-1}] - [Br_3^{-1}] - [Br^{-1}] +$$

$$[Na^{+1}] + [K^{+1}] - [HSO_4^{-1}] - 2[SO_4^{-2}] = 0 \quad (9a)$$

$$f_1 = f(H) :$$

$$2N_1 + N_2(1+2n_2) + N_3(1+2n_3) + N_4(1+2n_4) + 2N_5n_5 + N_6(1+2n_6) + 2N_7n_7 + 2N_8n_8 + 2N_9n_9 + 2N_{10}n_{10} + 2N_{11}n_{11} + 2N_{12}n_{12} + N_{13}(1+2n_{13}) + 2N_{14}n_{14} = 2N_{02} + 2N_{03} + 2N_{05}$$

$$f_2 = f(O) :$$

$$N_1 + N_2(1+n_2) + N_3(1+n_3) + N_4(3+n_4) +$$

$$N_5(3+n_5) + N_6(1+n_6) + N_7(1+n_7) + N_8n_8 +$$

$$N_9n_9 + N_{10}n_{10} + N_{11}n_{11} + N_{12}n_{12} + N_{13}(4+n_{13}) +$$

$$N_{14}(4+n_{14}) = 4N_{02} + N_{03} + 3N_{04} + N_{05}$$

$$-f_3 = -f(Na) :$$

$$N_{01} = N_{11} \Rightarrow [Na^{+1}] = C_0V_0/(V_0+V) \quad (10)$$

$$-f_4 = -f(K) :$$

$$N_{04} = N_{12} \Rightarrow [K^{+1}] = CV/(V_0+V) \quad (11)$$

$$-f_5 = -f(SO_4) \Leftrightarrow -f_5 = -f(S) : \quad (12)$$

$$N_{02} = N_{13} + N_{14} \Rightarrow \quad (12a)$$

$$[HSO_4^{-1}] + [SO_4^{-2}] = C_0V_0/(V_0+V) \quad (12b)$$

$$f_6 = f(Br) : N_4 + N_5 + N_6 + N_7 + 2N_8 + 3N_9 + N_{10}$$

$$= N_{01} + N_{04} \quad \Rightarrow \quad (13)$$

$$[HBrO_3] + [BrO_3^{-1}] + [HBrO] + [BrO^{-1}] + 2[Br_2] + 3[Br_3^{-1}] + [Br^{-1}] = (C_0V_0 + CV)/(V_0+V) \quad (13a)$$

$$f_{12} = 2 f(O) - f(H)$$

$$-N_2 + N_3 + 5N_4 + 6N_5 + N_6 + 2N_7 + 7N_{13} + 8N_{14} = 6N_{02} + 6N_{04} \quad (14)$$

$$- [H^{+1}] + [OH^{-1}] + 5[HBrO_3] + 6[BrO_3^{-1}] + [HBrO] + 2[BrO^{-1}] + 7[HSO_4^{-1}] + 8[SO_4^{-2}] = 6(C_0V_0 + CV)/(V_0+V) \quad (14a)$$

$$f_{12} + f_0 - f_3 - f_4 - 6 \cdot f_5 = 0 \quad \Leftrightarrow \quad (+1)f_1 +$$

$$(-2)f_2 + (+1)f_3 + (+1)f_4 + (+6)f_5 - f_0 = 0 \quad \Leftrightarrow$$

$$(+1)f(H) + (-2)f(O) + (+1)f(Na) + (+1)f(K) +$$

$$(+6)f(S) - \text{ChB} = 0 \quad (15)$$

$$5(N_4+N_5) + 1(N_6+N_7) - N_9 - N_{10}$$

$$= -N_{01} + 5N_{04} \quad \Rightarrow \quad (15a)$$

$$5([HBrO_3] + [BrO_3^{-1}]) + [HBrO] + [BrO^{-1}] -$$

$$[Br_3^{-1}] - [Br^{-1}] = -\frac{C_0V_0}{V_0+V} + 5 \cdot \frac{CV}{V_0+V} \quad (15b)$$

$$(+5)([HBrO_3]+[BrO_3^{-1}]) + (+1)([HBrO]+[BrO^{-1}]) + 2 \cdot 0 \cdot [Br_2] + 3 \cdot \left(-\frac{1}{2}\right) \cdot [Br_3^{-1}] + (-1)[Br^{-1}]$$

$$= (-1) \cdot \frac{C_0V_0}{V_0+V} + (+5) \cdot \frac{CV}{V_0+V} \quad (15c)$$

$$Z_{Br} \cdot f_6 - (f_{12} + f_0 - f_3 - f_4 - 6 \cdot f_5)$$

$$(Z_{Br}-5)(N_4+N_5) + (Z_{Br}-1)(N_6+N_7) + 2Z_{Br}N_8 + (3Z_{Br}+1)N_9 + (Z_{Br}+1)N_{10}$$

$$= (Z_{Br}+1)N_{01} + (Z_{Br}-5)N_{04} \quad \Rightarrow \quad (16)$$

$$(Z_{Br}-5)([HBrO_3]+[BrO_3^{-1}]) + (Z_{Br}-1)([HBrO]+[BrO^{-1}]) + 2Z_{Br}[Br_2] + (3Z_{Br}+1)[Br_3^{-1}] + (Z_{Br}+1)[Br^{-1}] =$$

$$(Z_{Br}+1) \cdot \frac{C_0V_0}{V_0+V} + (Z_{Br}-5) \cdot \frac{CV}{V_0+V} \quad (16a)$$

Other linear combinations are also possible. From the linear combination  $(f_{12} + f_0 - f_3 - f_4 - 6 \cdot f_5 + f_6)/2$  we get the shortest (involving the smallest number of components) form of GEB for the system S:

$$3(N_4+N_5) + (N_6+N_7) + N_8 + N_9 = 3N_{04} \quad (17)$$

$$3([HBrO_3] + [BrO_3^{-1}]) + ([HBrO] + [BrO^{-1}]) + [Br_2] + [Br_3^{-1}] = 3 \cdot \frac{CV}{V_0+V} \quad (17a)$$

Note that the coefficients in eq. 15c are equal to the oxidation numbers (ONs) of the elements considered as 'fans' in the system S, i.e., H, O, Na, K, S. Eq. 15b obtained from the combination 15 interrelates components and species formed by Br, as the player.

Other linear combinations  $a_1 \cdot f_0 + a_2 \cdot f_{12} + \sum_{k=3}^6 a_k \cdot f_k$  of the balances:  $f_0, f_{12}, f_3, \dots, f_6$  are also acceptable,  $a_k \in \mathbb{R}$ .

Equations 14a, 15b, 16a and 17a are alternative/equivalent equations for GEB related to the system S. One of the equations for GEB, together with equations 9a, 12b and 13a, form a complete set of equations related to the system S. The relations 10 and 11, considered as equalities (not equations), can be immediately introduced into eq. 9a as numbers.

In eq. 12b, the  $\text{SO}_4^{-2}$  can be perceived as the core. However, because the system S has no other competing sulfate forms, the choice between  $f(\text{S})$  and  $f(\text{SO}_4)$  (eq. 12) is irrelevant.

### Separate Approaches to D and T of the System S

We can also formulate the balances for D and T, considered separately, as independent units. Applying the notation specified above, we have:

- for D :

$$f_0 = \text{ChB} : N_2 - N_3 - N_{10} + N_{11} - N_{13} - 2N_{14} = 0$$

$$f_1 = f(\text{H}) : 2N_1 + N_2(1+2n_2) + N_3(1+2n_3) +$$

$$2N_{11}n_{11} + N_{13}(1+2n_{13}) + 2N_{14}n_{14} = 2N_{02} + 2N_{03}$$

$$f_2 = f(\text{O}) : N_1 + N_2n_2 + N_3(1+n_3) + N_{11}n_{11} +$$

$$N_{13}(4+n_{13}) + N_{14}(4+n_{14}) = 4N_{02} + N_{03}$$

$$-f_3 = -f(\text{Na}) : N_{01} = N_{11}$$

$$-f_5 = -f(\text{SO}_4) : N_{02} = N_{13} + N_{14}$$

$$f_6 = f(\text{Br}) : N_{10} = N_{01}$$

and then:

$$f_{12} = 2 \cdot f_2 - f_1 : -N_2 + N_3 + 7N_{13} + 8N_{14} = 6N_{02}$$

$$f_{12} + f_0 - f_3 - 6f_5 + f_6 = 0 \Leftrightarrow (+1)f(\text{H}) + (-2)f(\text{O}) + (+1)f(\text{Na}) + (+6)f(\text{S}) + (-1)f(\text{Br}) - \text{ChB} = 0 \quad (18)$$

$$0 = 0 \quad (18a)$$

- for T :

$$f_0 = \text{ChB} : N_2 - N_3 - N_5 + N_{12} = 0$$

$$f_1 = f(\text{H}) : 2N_1 + N_2(1+2n_2) + N_3(1+2n_3) +$$

$$N_4(1+2n_4) + 2N_5n_5 + 2N_{12}n_{12} = 2N_{05}$$

$$f_2 = f(\text{O}) : N_1 + N_2n_2 + N_3(1+n_3) + N_4(3+n_4) +$$

$$N_5(3+n_5) + N_{12}n_{12} = 3N_{04} + N_{05}$$

$$-f_4 = -f(\text{K}) : N_{04} = N_{12}$$

$$-f_6 = -f(\text{Br}) : N_{04} = N_4 + N_5$$

and then:

$$f_{12} = 2 \cdot f_2 - f_1 : -N_2 + N_3 + 5N_4 + 6N_5 + 8N_{14} = 6N_{04}$$

$$f_{12} + f_0 - f_4 - 5f_6 = 0 \Leftrightarrow (+1)f(\text{H}) + (-2)f(\text{O}) + (+1)f(\text{K}) + (+5)f(\text{Br}) - \text{ChB} = 0 \quad (19)$$

$$0 = 0 \quad (19a)$$

The D and T, considered separately, form non-redox-systems; ONs for Br are:  $-1$  in NaBr, and  $+5$  in  $\text{KBrO}_3$ , i.e., there are the boundary values of ONs in

bromine redox systems; Br(+7) species are omitted in considerations [24].

The relations  $0 = 0$ , named as identities, mean here that:

- $f_{12}$  is linearly dependent on:  $f_0, f_3, f_5$  and  $f_6$  in eq. 18, i.e.,  $f_{12} = f_3 + 6f_5 - f_6 - f_0$ ;
- $f_{12}$  is linearly dependent on:  $f_0, f_4$  and  $f_6$  in eq. 19, i.e.,  $f_{12} = f_4 + 5f_6 - f_0$ .

In other words, the  $f_{12}$  are not the independent equations in D and T, considered here as separate subsystems.

### Approach I to GEB in the System S

Unlike the Approach II exemplified above, the Approach I to GEB needs prior knowledge of ONs for all elements in components and species of the system in question. In the system S, there are  $K^* = 5$  'fans', whereas bromine (Br) is considered as the 'player',  $K - K^* = 6 - 5 = 1$  is here the number of players.

In the system S, bromine (as NaBr and  $\text{KBrO}_3$ ) is the carrier/distributor of the player electrons. One atom of Br has  $Z_{\text{Br}}$  bromine electrons, and then one molecule of NaBr has  $Z_{\text{Br}} + 1$  bromine electrons, one molecule of  $\text{KBrO}_3$  has  $Z_{\text{Br}} - 5$  bromine electrons; then  $N_{01}$  molecules of NaBr involve  $(Z_{\text{Br}} + 1) \cdot N_{01}$  bromine electrons,  $N_{04}$  molecules of  $\text{KBrO}_3$  involve  $(Z_{\text{Br}} - 5)N_{04}$  bromine electrons. Thus, the total number of bromine electrons introduced by NaBr and  $\text{KBrO}_3$  is  $(Z_{\text{Br}} + 1) \cdot N_{01} + (Z_{\text{Br}} - 5)N_{04}$ . On this basis, we state that [2,27]:

$N_4$  species  $\text{HBrO}_3 \cdot n_4 \text{H}_2\text{O}$  involve  $(Z_{\text{Br}} - 5) \cdot N_4$  bromine electrons;

$N_5$  species  $\text{BrO}_3^{-1} \cdot n_5 \text{H}_2\text{O}$  involve  $(Z_{\text{Br}} - 5) \cdot N_5$  bromine electrons;

$N_6$  species  $\text{HBrO} \cdot n_6 \text{H}_2\text{O}$  involve  $(Z_{\text{Br}} - 1) \cdot N_6$  bromine electrons;

$N_7$  species  $\text{BrO}^{-1} \cdot n_7 \text{H}_2\text{O}$  involve  $(Z_{\text{Br}} - 1) \cdot N_7$  bromine electrons;

$N_8$  species  $\text{Br}_2 \cdot n_8 \text{H}_2\text{O}$  involve  $2Z_{\text{Br}} \cdot N_8$  bromine electrons;

$N_9$  species  $\text{Br}_3^{-1} \cdot n_9 \text{H}_2\text{O}$  involve  $(3Z_{\text{Br}} + 1) \cdot N_9$  bromine electrons;

$N_{10}$  species  $\text{Br}^{-1} \cdot n_{10} \text{H}_2\text{O}$  involve  $(Z_{\text{Br}} + 1) \cdot N_{10}$  bromine

electrons.

The balance for the bromine electrons is then expressed by eq. 16 and then by eq. 16a. This confirms the equivalency of the Approaches I and II to GEB (eq. 1).

As stated above, the Approach I to GEB is compared to the 'card game' [27] (pp. 41-43), and – nominally – all electrons of the players are involved in the balance 16a. Following this line of reasoning, it can be also stated that the card players do not engage, as a rule, all their cash resources in the game. What's more – the 'debt of honour' principle can be applied [27] (p. 43). Simply, on the ground of linear combination, in eq. 16a one can replace  $Z_{Br}$  for Br by  $\zeta_{Br} \neq Z_{Br}$ ; in particular, one can apply  $\zeta_{Br} = 0$ , see equations 15a and 15b in context with equations 16 and 16a.

### Computer Program for the System S

function F = Function\_NaBr\_H2SO4\_KBrO3(x)

%NaBr (C0) H2SO4 (C01) V0 BkBrO3 (C) V

global V Vmin Vstep Vmax V0 C C0 C01 fi H OH pH E Kw  
pKw A

global Br Br2 Br3 HBrO BrO HBrO3 BrO3 Na K HSO4 SO4

global logBr logBr2 logBr3 logHBrO logBrO logHBrO3  
logBrO3

global logNa logK logHSO4 logSO4

E=x(1);

pH=x(2);

Br=10.^-x(3);

SO4=10.^-x(4);

H=10.^-pH;

pKw=14;

Kw=10.^-14;

OH=Kw./H;

A=16.9;

ZBr=35;

BrO3=Br.\*10.^(6.\*A.\*(E-1.45)+6.\*pH);

HBrO3=10.^0.7.\*H.\*BrO3;

BrO=Br.\*10.^(2.\*A.\*(E-0.76)+2.\*pH-2.\*pKw);

HBrO=10.^8.6.\*H.\*BrO;

Br3=Br.^3.\*10.^(2.\*A.\*(E-1.05));

Br2=Br.^2.\*10.^(2.\*A.\*(E-1.087));

Na=C0.\*V0./(V0+V);

K=C.\*V./(V0+V);

%Charge balance

F=[(H-OH-BrO3-BrO-Br3-Br+Na+K-HSO4-2.\*SO4);

%Concentration balance of Br

(HBrO3+BrO3+HBrO+BrO+2.\*Br2+3.\*Br3+

Br-(C0.\*V0.+C.\*V)/(V0+V));

%Concentration balance for SO4

(HSO4+2.\*SO4-C01.\*V0)/(V0+V));

%GEB

((ZBr-5).\*(HBrO3+BrO3)+(ZBr-1).\*(HBrO+BrO)

+2.\*ZBr.\*Br2...

+(3.\*ZBr+1).\*Br3+(ZBr+1).\*Br...

-((ZBr+1).\*C0.\*V0+(ZBr-5).\*C.\*V)/(V0+V));

logBr=log10(Br);

logBr2=log10(Br2);

logBr3=log10(Br3);

logHBrO=log10(HBrO);

logBrO=log10(BrO);

logHBrO3=log10(HBrO3);

logBrO3=log10(BrO3);

logHSO4=log10(HSO4);

logSO4=log10(SO4);

logNa=log10(Na);

logK=log10(K);

### Graphical Presentation of Results for the System S

In the algorithm, prepared according to MATLAB

computational software, the potential  $E$  [V] was expressed in SHE scale [46],  $\text{pH} = -\log[\text{H}^{+1}]$ ,  $\text{pBr} = -\log[\text{Br}^{-1}]$ ,  $V_0 = 100$ ,  $C_0 = 0.01$ ,  $C = 0.1$ ,  $C_{01} = 0$ ,  $0.01$  or  $0.1$ . The equilibrium constants related to this system were cited in ref. [2].


The results of calculations are presented graphically (Fig. 1), as the graphs: (a)  $E = E(\Phi)$ , (b)  $\text{pH} = \text{pH}(\Phi)$  and (c) speciation curves  $[X_i^{z_i}] = \varphi_i(\Phi)$ , with the fraction titrated  $\Phi$  (eq. 8) on the abscissas. The impact of growth in  $C_{01}$  concentration is illustrated here.

At  $C_{01} = 0$ , comproportionation practically does not occur (IIIa); concentration of  $\text{HBrO}$ , as the major product formed in the comproportionation reaction



is ca.  $10^{-6}$  mol/L. The potential  $E$  increases monotonically (Fig. 1a), whereas  $\text{pH}$  first increases, passes through maximum and then decreases, see IIa). The relevant  $E$  and  $\text{pH}$  changes are small (Figures 1a, IIa). Binding the  $\text{H}^{+1}$  ions in reaction 21 causes a weakly alkaline reaction (Fig. IIa).

At  $C_{01} = 0.01$  (Figures 1b, IIb, IIIb) and  $0.1$  (Figures 1c, IIc, IIIc), the stoichiometry  $1 : 5$ , i.e.,  $\Phi_{\text{eq}} = 0.2$ , see [16,28], resulting from the shape of the related graphs, is expressed by reaction



For  $\Phi > 0.2$ , an increase of efficiency of the competing reaction 21 is noted. A growth of  $C_{01}$  value causes a small extension of the potential range in the jump region, on the side of higher  $E$ -values (Fig. 1, column I). With an increase of the  $C_{01}$  value, the graphs of  $\text{pH}$  vs.  $\Phi$  resemble two almost straight line segments intersecting at  $\Phi_{\text{eq}} = 0.2$  (Fig. 2, column II). However, the  $\text{pH}$ -ranges covered by the titration curves are gradually narrowed; it is an effect in growth of dynamic buffer capacity of the related redox systems [32].

### Concluding remarks

The linear combination

$$f_0 + f_{12} - \sum_{k=3}^K d_k \cdot f_k = 0 \quad \Leftrightarrow$$

$$\sum_{k=3}^K d_k \cdot f_k - (f_{12} + f_0) = 0 \quad \Leftrightarrow$$

$$f_1 - 2 \cdot f_2 + \sum_{k=3}^K d_k \cdot f_k - f_0 = 0 \quad \Leftrightarrow$$

$$\sum_{k=1}^K d_k \cdot f_k - f_0 = 0 \quad (22)$$

involves  $K$  balances:  $f_0, f_{12}, f_3, \dots, f_K$  where  $d_1 = +1$ ,  $d_2 = -2$ . All the balances are presented here as equations,  $f_k = 0$ .

In a non-redox system, we have  $K$  fans, i.e., the number of players equals zero. In a redox system, we have  $K^*$  fans,  $K^* < K$ , i.e., the number of players equals  $K - K^* (> 0)$ .

When the multipliers  $d_k$  are equal to (or involved with) the oxidation numbers (ONs) of the corresponding elements ( $k = 1, \dots, K$ ) in a non-redox system, then eq. 22 is transformed into identity,  $0 = 0$ . This proves that  $f_{12}$  is not the independent equation in the set  $f_0, f_{12}, f_3, \dots, f_K$ , and  $f_{12} = \sum_{k=3}^K d_k \cdot f_k - f_0$ . Then  $f_0, f_3, \dots, f_K$  is the set of  $K-1$  independent balances, composed of charge balance ( $f_0$ ) and  $K-2$  elemental/core balances  $f_3, \dots, f_K$ .

Referring now to a redox system, we arrange the elemental/core balances in the sequence  $f_1, f_2, \dots, f_{k^*}, f_{k^*+1}, \dots, f_K$ , and then formulate the equation

$$\sum_{k=1}^{K^*} d_k \cdot f_k - f_0 = 0 \quad (23)$$

involving the balances for  $K^*$  electron-non-active elements, compare with equations 15, 18, 19.

If  $d_k$  ( $k=1, \dots, K^*$ ) are equal to (or involved with) ONs of electron-non-active elements, then the resulting balance involves only the components ( $N_i, N_{0j}$ ) related to electron-active species/compounds, with coefficients equal to (or involved with) ONs of these element. More precisely,  $d_k$  are the products of the number  $\lambda_k$  of defined atoms in the species and the ON value  $\omega_k$ ,  $d_k = \lambda_k \cdot \omega_k$  i.e. ( $k = 1, \dots, K^*$ ). It is clearly visible in the case of  $\text{Br}_3^{-1}$  in eq. 15c, where  $-1 = 3 \cdot (-\frac{1}{3})$ , and (less visible) in the case of  $\text{Br}_2$  in the eq. 15c, where  $0 = 2 \cdot 0$ , i.e., the coefficient  $d_9 = -1$  at  $[\text{Br}_3^{-1}]$  in eq. 15b is involved with (not equal to) the oxidation number  $\omega_9 = (-\frac{1}{3})$  of bromine in  $\text{Br}_3^{-1}$  ( $\lambda_9 = 3$ ).

The equivalent relations were applied:

$$f_k = \sum_{i=1}^I a_{ki} \cdot N_i - \sum_{j=1}^J b_{kj} \cdot N_{0j} = 0 \Leftrightarrow \sum_{i=1}^I a_{ki} \cdot N_i = \sum_{j=1}^J b_{kj} \cdot N_{0j} \quad (24)$$

for elements with negative oxidation numbers, or

$$-f_k = \sum_{j=1}^J b_{kj} \cdot N_{0j} - \sum_{i=1}^I a_{ki} \cdot N_i = 0 \quad \Leftrightarrow$$

$$\sum_{j=1}^I b_{kj} \cdot N_{0j} = \sum_{i=1}^I a_{ki} \cdot N_i \quad (25)$$

for elements with positive oxidation numbers,  $k \in 3, \dots, K$ . The change of places of numbers  $N_i$  for components and  $N_{0j}$  for species at the equality sign in relations 24, 25 were made in order to avoid possible/simple mistakes in the realization of the linear combination procedure. This facilitates the purposeful linear combination of the balances, and enables to avoid simple mistakes in this operation. Note, for example, that  $f_4 = f(\text{Na}) \Leftrightarrow f(\text{Na}) = f_4 \Leftrightarrow -f_4 = -f(\text{Na})$ .

Starting from  $K+1$  balances:  $f_0, f_1, f_2, f_3, \dots, f_k$ , after formulation of the linear combination  $f_{12}$  we obtain the set of  $K$  balances:  $f_0, f_{12}, f_3, \dots, f_k$ . In a non-redox system,  $f_{12}$  is the dependent balance; we have there  $K-1$  independent balances:  $f_0, f_3, \dots, f_k$ . In other words,  $f_{12}$  is not a new, independent balance in non-redox systems; it is then omitted in formulation of any non-redox system. The identity  $0 = 0$  for the linear combinations indicates that the equations  $f_0, f_{12}, f_3, \dots, f_k$  are linearly dependent for non-redox systems. In a redox system,  $f_{12}$  is the independent balance (i.e., different from the identity,  $0=0$ ); then we have  $K$  independent balances:  $f_0, f_{12}, f_3, \dots, f_k$ , that will be rearranged – optionally – as the set  $(f_{12}, f_0, f_3, \dots, f_k)$  involved with GEB, ChB, and  $f(Y_k)$  ( $k = 3, \dots, K$ ), respectively. The number of elemental/core balances, both in non-redox and redox systems, equals  $K-2$ . Then:

- For a non-redox system, a proper linear combination of  $f_{12}$  with  $f_0, f_3, \dots, f_k$  is reducible to identity  $0 = 0$ , i.e.,  $f_{12}$  is linearly dependent on  $f_0, f_3, \dots, f_k$  (equations 24,25).
- For a redox system, any linear combination of  $f_{12}$  with  $f_0, f_3, \dots, f_k$  is not reducible to identity  $0 = 0$ , i.e.,  $f_{12}$  is linearly independent on  $f_0, f_3, \dots, f_k$ .
- In conclusion, the linear independency/dependency of  $f_{12} = 2 \cdot f(\text{O}) - f(\text{H})$  from other balances:  $f_0, f_3, \dots, f_k$  is the general criterion distinguishing between redox and non-redox systems; the proper linear combination LC with  $d_k$  equal to ON's (see eq. 15), is the way towards the simplest/shortest form of GEB; the shortest form (eq. 17a) was obtained after further combination with the balance for Br (player). For a non-redox system, the linear combination indicated it is the way towards identity  $0 = 0$  (equations 18a, 19a).
- $f_{12} = 2 \cdot f(\text{O}) - f(\text{H})$  is the *primary* form of Generalized Electron Balance (GEB),  $f_{12} = pr$ GEB, completing the set

of  $K$  balances  $f_0, f_{12}, f_3, \dots, f_k$  necessary for resolution of redox systems of any degree of complexity.

- Any linear combination of  $f_{12}$  with the balances  $f_0, f_3, \dots, f_k$  has full properties of GEB related to the redox system considered, i.e., all them are equivalent forms of GEB.
- The Approach II to GEB does not indicate oxidants and reductants, i.e., oxidized and reduced forms in the system in question.
- The prior knowledge of oxidation numbers (ONs) for all elements of the system is not required; this fact is of capital importance when redox equilibria are involved, e.g., with complex organic species; the known composition of a species, expressed by its formula, together with external charge of this species, provides information sufficient to formulate the related balances.
- When the oxidation numbers of all elements of a system are known beforehand, the GEB can be formulated according to Approach I to GEB, known also as the 'short version' of GEB; the GEB obtained according to Approach I to GEB involves all electron-active elements – as components and species – of the system tested.
- Both Approaches (I and II) to GEB are equivalent (eq. 1).

where all 'fans' ( $K^*$  elements or cores) of the system are involved;  $K^* = K$  for a non-redox system,  $K^* < K$  for a redox system, where  $K-K^*$  players are involved. When  $d_k$  are equal to the oxidation numbers (ONs) of elements in the corresponding 'fans' ( $k=1, \dots, K$ ) of a non-redox system, then LC is transformed into identity,  $0 = 0$ . For a redox system, LC assumes there simpler/simplest form, where only the species and components related to players are involved. The  $f_{12}$  is considered as the *primary* form of GEB,  $f_{12} = pr$ -GEB. The  $f_{12}$ , LC and any other combination

$$\sum_{k=3}^K a_k \cdot f_k - (f_{12} + a_0 \cdot f_0),$$

where  $a_k \in \mathbb{R}$ , have full properties of GEB, although the simplest/shortest form of GEB, involving the smallest number of components, is more desirable.

- The criterion distinguishing between non-redox and redox systems is valid for redox systems of any degree of complexity. We can check it also on more complex redox systems, of any degree of complexity, where







Fig. 1: The functions:  $E = E(\Phi)$  (column I),  $\text{pH} = \text{pH}(\Phi)$  (column II) and speciation diagrams (column III), at  $(C_0, V_0, C) = (0.01, 100, 0.1)$ , and three different  $C_{01}$  values, indicated at the graphs.

two or more electron-active elements as 'players', are involved. The relation (12) was also confirmed for electrolytic systems in binary and (generally) mixed-solvent  $A_s$  ( $s = 1, \dots, S$ ) media [47-49], where mixed solvates



are assumed, and  $n_{iA_s} \geq 0$  is the mean numbers of  $A_s$  molecules attached to  $X_i^{z_i}$ .

- The formulation of GEB according to Approach II is relatively/extraordinarily simple, although receiving the shorter equation for GEB, when using the linear combinations of *pr*-GEB with other balances, requires implementation of the time-consuming, preparatory activities. However, the formulation of GEB according to Approach II has – undoubtedly – the cognitive advantages, even in the cases when the oxidation numbers for all elements in the system are known beforehand.

### Final comments

The GEB concept, discovered by Michałowski as the Approaches I (1992, [2-4]) and II (2005, [8]) to GEB, is perceived as the law of the matter conservation, as the general Law of Nature [27], as the hidden connection of physicochemical laws, and as a breakthrough in thermodynamic theory of electrolytic redox systems. The GATES/GEB formulated in 1992, is an example of excellent paradigm, as the best possible thermodynamic approach to electrolytic redox systems. Other/earlier approaches to formulation of electrolytic redox systems, based on stoichiometric principles, were unsuccessful, for obvious reasons. All them were thoroughly criticized in our review papers [15-17, 25-30], and elsewhere.

The Approach II to GEB does not require any prior knowledge of the oxidation numbers of elements in the components and in the species; it is a fact of capital importance, particularly in relation to organic species (molecules, ions, radicals and ion-radicals), of any degree of complexity. Note that the 'oxidation number' was essentially the contractual concept [50-52], here it results naturally from the linear combination; within the Approach II to GEB, ON is then the derivative concept. If the oxidation numbers are easily determined, the Approach I to GEB, known as the 'short' version of GEB, can be applied. Within the Approaches I and II to GEB,

the roles/terms of oxidants and reductants are not assigned *a priori* to individual components; GATES/GEB provides full 'democracy' in this regard.

All the inferences made within GATES/GEB are based on firmly established physical, physicochemical and algebraic foundations. Consequently, it allows to understand far better the physicochemical phenomena occurring in the system in question, and improve some methods of analysis. All the facts testify very well about the potency of simulated calculations made, according to GATES, on the basis of all attainable and preselected physicochemical knowledge involved.

Contrary to appearances, established by the current paradigm, the criterion distinguishing non-redox and redox systems is not immediately associated with free electrons in the related system; it provides a kind of uniformity in the formulas derived for this purpose. This fact, especially the simple calculations ([26], pp. 361-363) of free electrons concentrations in redox systems, deny the unique role of free electrons in elementary redox reactions, as described in half- or partial reactions, where the species of the same element with different oxidation numbers are involved. On the other hand, it points to the unique role of H and O in redox systems, suggested in earlier theoretical/hypothetical/qualitative considerations on these systems. Let's repeat: the GEB concept, valid for redox electrolytic systems, is the emanation of balances for H and O, referred to aqueous and non-aqueous media, with amphiprotic (co)solvents involved.

### Notations and Acronyms

D – titrand (solution titrated),

GATES – Generalized Approach to Electrolytic Systems,

GEB – Generalized Electron Balance,

ON – oxidation number,

T – titrant,

V – volume (mL) of T,

V0 – volume (mL) of D.

### References

1. Marx D., Tuckerman M.E., Hutter J., Parrinello M. (1999) The nature of the hydrated excess proton in water, *Nature*, 397: 601-604.

2. Michałowski T. (1994) Calculation of pH and potential E for bromine aqueous solutions, *Journal of Chemical Education*, 71(7): 560-562.
3. Michałowski T., Lesiak A. (1994) Acid-base titration curves in disproportionating redox systems, *Journal of Chemical Education* 71(8): 632-636.
4. Michałowski T., Lesiak A. (1994) Formulation of generalized equations for redox titration curves, *Chemia Analityczna (Warsaw)*, 39: 623-637.
5. Michałowski T., Wajda N., Janecki D. (1996) A Unified Quantitative Approach to Electrolytic Systems, *Chemia Analityczna (Warsaw)*. 41: 667-685.
6. Michałowski T., Baterowicz A., Madej A., Kochana J. (2001) An extended Gran method and its applicability for simultaneous determination of Fe(II) and Fe(III), *Analytica Chimica Acta*. 442(2): 287-293.
7. Michałowski T., Toporek M., Rymanowski M. (2005) Overview on the Gran and other linearization methods applied in titrimetric analyses, *Talanta* 65 (5): 1241-1253.
8. Michałowski T., Pietrzyk A. (2006) The generalized electron balance concept. Derivation based on elementary rules of the matter conservation, in: *Chemometrics: Methods and Applications* (ed. D. Zuba, A. Parczewski), Kraków. 415-422.
9. Michałowski T., Rymanowski M., Pietrzyk A. (2005) Non-typical Brönsted's acids and bases, *Journal of Chemical Education*. 82(3): 470-472.
10. Michałowski T. (2007) Complementarity of physical and chemical laws of preservation in aspect of electrolytic systems (in Polish), *Wiadomości Chemiczne*. 61: 625-640.
11. Michałowski T. (2007) Complementarity of physical and chemical laws of preservation in aspect of electrolytic systems (in Polish), *Use of information technology in academic teaching of chemistry*, UJ Cracow. 155-161.
12. Michałowski T. (2007) Electron balance as the basis of calculations in redox systems (in Polish), *Use of information technology in academic teaching of chemistry*, UJ Cracow. 162-169.
13. Michałowski T, Kupiec K, Rymanowski M. (2008) Numerical analysis of the Gran methods. A comparative study, *Analytica Chimica Acta* 606: 172-183.
14. Ponikvar M., Michałowski T., Kupiec K., Wybraniec S., Rymanowski M. (2008) Experimental verification of the modified Gran methods applicable to redox systems, *Analytica Chimica Acta* 628: 181-189.
15. Michałowski T. (2010) The Generalized Approach to Electrolytic Systems: I. Physicochemical and Analytical Implications, *Critical Reviews in Analytical Chemistry*, 40: 2-16.
16. Michałowski T., Pietrzyk A., Ponikvar-Svet M., Rymanowski M. (2010) The Generalized Approach to Electrolytic Systems: II. The Generalized Equivalent Mass (GEM) Concept, *Critical Reviews in Analytical Chemistry* 40: 17-29.
17. Michałowski T. (2011) Application of GATES and MATLAB for Resolution of Equilibrium, Metastable and Non-Equilibrium Electrolytic Systems, Chapter 1 (p. 1-34) in: *Applications of MATLAB in Science and Engineering* (ed. by T. Michałowski), InTech-Open Access publisher in the fields of Science, Technology and Medicine.
18. Michałowski T., Ponikvar-Svet M., Asuero A.G., Kupiec K. (2012) Thermodynamic and kinetic effects involved with pH titration of As(III) with iodine in a buffered malonate system, *Journal of Solution Chemistry* 41(3): 436-446.
19. Michałowska-Kaczmarczyk, A.M., Michałowski, T. (2013) Comparative balancing of non-redox and redox electrolytic systems and its consequences, *American Journal of Analytical Chemistry*. 4: 46-53.
20. Michałowska-Kaczmarczyk A.M., Michałowski T. (2014) GATES as the Unique Tool for Simulation of Electrolytic Redox and Non-Redox Systems, *Journal of Analytical & Bioanalytical Techniques* 5(4): 1-5.
21. Michałowski T., Toporek M., Michałowska-Kaczmarczyk A.M., Asuero A.G. (2013) New Trends in Studies on Electrolytic Redox Systems. *Electrochimica Acta* 109: 519-531.
22. Michałowski T., Michałowska-Kaczmarczyk A.M., Toporek M. (2013) Formulation of general criterion

- distinguishing between non-redox and redox systems, *Electrochimica Acta* 112: 199-211.
23. Michałowska-Kaczmarczyk A.M., Toporek M., Michałowski T. (2015) Speciation Diagrams in Dynamic Iodide + Dichromate System, *Electrochimica Acta* 155: 217–227.
  24. Toporek M., Michałowska-Kaczmarczyk A.M., Michałowski T. (2015) Symproportionation versus Disproportionation in Bromine Redox Systems, *Electrochimica Acta* 171: 176–187.
  25. Michałowska-Kaczmarczyk A.M., Asuero A.G., Toporek M., Michałowski T., “Why not stoichiometry” versus “Stoichiometry – why not?” Part II. GATES in context with redox systems, *Critical Reviews in Analytical Chemistry* 45(3): (2015) 240–268.
  26. Michałowska-Kaczmarczyk A.M., Michałowski T., Toporek M., Asuero A.G. (2015) “Why not stoichiometry” versus “Stoichiometry – why not?” Part III, Extension of GATES/GEB on Complex Dynamic Redox Systems, *Critical Reviews in Analytical Chemistry*, 45(4): 348-366. DOI: 10.1080/10408347.2014.953673.
  27. Michałowska-Kaczmarczyk A.M., Spórna-Kucab A., Michałowski T. (2017) Generalized Electron Balance (GEB) as the Law of Nature in Electrolytic Redox Systems, in: *Redox: Principles and Advanced Applications*, Ali Khalid, M.A. (Ed.) InTech, Chap. 2, 9-55, DOI: 10.5772/66005 (248 pages).
  28. Michałowska-Kaczmarczyk A.M., Spórna-Kucab A., Michałowski T. (2017) Principles of Titrimetric Analyses According to Generalized Approach to Electrolytic Systems (GATES), in: *Advances in Titration Techniques*, Vu Dang Hoang (Ed.) InTech, Chap 5: 133-171, DOI: 10.5772/66980 (216 pages).
  29. Michałowska-Kaczmarczyk A.M., Spórna-Kucab A., Michałowski T. (2017) A Distinguishing Feature of the Balance  $2 \cdot f(O) - f(H)$  in Electrolytic Systems. The Reference to Titrimetric Methods of Analysis, in: *Advances in Titration Techniques*. Vu Dang Hoang (Ed.) InTech, Chap 6: 173-207.
  30. Michałowska-Kaczmarczyk A.M., Spórna-Kucab A., Michałowski T. (2017) Solubility products and solubility concepts, in: *Descriptive Inorganic Chemistry. Researches of Metal Compounds*. Akitsu T. (Ed.) InTech, Chap 5: 93-134.
  31. Michałowska-Kaczmarczyk A.M., Spórna-Kucab A., Michałowski T. (2017) Formulation of simple electrolytic redox systems according to GATES/GEB principles, *Journal of Chemistry and Applied Chemical Engineering*, 1(1): 1-10.
  32. Michałowska-Kaczmarczyk A.M., Spórna-Kucab A., Michałowski T. (2017) Dynamic Buffer Capacities in Redox Systems, *Journal of Chemistry and Applied Chemical Engineering*, 1(2): 1–7.
  33. Michałowska-Kaczmarczyk A.M., Michałowski T., Toporek M. (2016) Formulation of Dynamic Redox Systems according to GATES/GEB Principles, *International Journal of Electrochemical Science*, 11: 2560-2578.
  34. Meija J., Michałowska-Kaczmarczyk A.M., Michałowski T. (2017) Redox titration challenge, *Analytical and Bioanalytical Chemistry*. 409(1): 11–13; DOI: 10.1007/s00216-016-0020-0.
  35. Michałowski T., Michałowska-Kaczmarczyk A.M., Meija J. (2017) Solution of redox titration challenge, *Analytical and Bioanalytical Chemistry*. 409(17): 4113-4115.
  36. Michałowska-Kaczmarczyk A.M., Spórna-Kucab A., Michałowski T. (2017) Some Regularities Involved with Oxidation Numbers Stated in Formulation of Redox Systems According to GATES/GEB Principles, *Journal of Analytical, Bioanalytical and Separation Techniques*. 2(2): 102-110.
  37. Michałowska-Kaczmarczyk A.M., Spórna-Kucab A., Michałowski T. (2018) The balance  $2 \cdot f(O) - f(H)$  as a keystone in formulation of electrolytic systems, *Research and Reviews in Computational Chemistry*. 1(1): 1-9.
  38. Michałowska-Kaczmarczyk A.M., Spórna-Kucab A., Michałowski T. (2017) General Properties of the Balances  $2 \cdot f(O) - f(H)$  Related to Electrolytic Systems, *Analytical Chemistry: An Indian Journal*, 17 (2): 124-.
  39. Michałowska-Kaczmarczyk A.M., Spórna-Kucab A., Michałowski T. (2017) A general property differentiating between redox and non-redox electrolytic systems and its consequences,

- International Journal of Mathematics and Statistics Invention 6(3): 67-76.
40. Michałowska-Kaczmarczyk A.M., Michałowski T. (2018) The importance of linear algebra in theory of electrolytic systems, Austin Chemical Engineering. 5 (1) in print.
41. Linear combination.
42. Emmy Noether.
43. Moore G.H. (1995) The axiomatization of linear algebra: 1875-1940, Historia Mathematica 22(3): 262-303.
44. Comproportionation.
45. Asuero A.G., Michałowski, T. (2011) Comprehensive formulation of titration curves referred to complex acid-base systems and its analytical implications, Critical Reviews in Analytical Chemistry 41(2): 151-187.
46. Standard hydrogen electrode
47. Michałowski T., Pilarski B., Asuero A.G., Michałowska-Kaczmarczyk A.M. (2014) Modeling of Acid-Base Properties in Binary-Solvent Systems, Chapter 9.4, pp. 623-648 in "Handbook of Solvents", Vol 1 Properties, Wypych G. (Editor), ChemTec Publishing, Toronto.
48. Michałowska-Kaczmarczyk A.M., Michałowski T. (2014) Generalized Electron Balance for Dynamic Redox Systems in Mixed-Solvent Media, Journal of Analytical Sciences, Methods and Instrumentation 4 (4): 102-109.
49. Michałowska-Kaczmarczyk A.M., Michałowski T. (2014) Linear Dependence of Balances for Non-Redox Electrolytic Systems, American Journal of Analytical Chemistry 5: 1285-1289.
50. Karen P., McArdle P., Takats J. (2016) Comprehensive definition of oxidation state (IUPAC Recommendations 2016). Pure and Applied Chemistry. 88(10): 831-839.
51. Karen P. (2015) Oxidation state, a long-standing issue! Angewandte Chemie International Edition. 54 (16): 4716-4726.
52. Karen P, McArdle P, Takats J. (2014) Toward a comprehensive definition of oxidation state (IUPAC Technical Report). Pure and Applied Chemistry. 86 (6): 1017-1081.